

de Sitter inflationary expansion from a noncompact KK theory: a nonperturbative quantum (scalar) field formalism

¹Mauricio Bellini*

¹*Departamento de Física, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata and Consejo Nacional de Ciencia y Tecnología (CONICET), Funes 3350, (7600) Mar del Plata, Argentina.*

Abstract

We develop a nonperturbative quantum scalar field formalism from a non-compact Kaluza-Klein (KK) theory using the induced-matter theory of gravity during inflation. We study the particular case of a de Sitter expansion for the universe.

Pacs numbers: 04.20.Jb, 11.10.kk, 98.80.Cq

I. INTRODUCTION

The two current versions of 5D gravity theory are membrane theory [1–3] and induced-matter theory [4]. In the former, gravity propagates freely into the bulk, while the interactions of particle physics are confined to a hypersurface (the brane). The induced-matter theory in its simplest form is the basic Kaluza-Klein (KK) theory in which the fifth dimension is not compactified and the field equations of general relativity in 4D follow from the fact that the 5D manifold is Ricci-flat; the large extra dimension is thus responsible for the appearance of sources in 4D general relativity. Hence, the 4D world of general relativity is embedded in a 5D Ricci-flat manifold. An interesting result of the induced-matter theory is that if $ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu$ is the 4D metric of any matter-free spacetime in general relativity, the $dS^2 = (\psi/\psi_0)^2 ds^2 - d\psi^2$ is the metric of a 5D manifold that is Ricci-flat [5,6].

During the last two decades the inflationary paradigm has become an almost universally accepted scenario to explain the observed large scale flatness and homogeneity of the universe [7]. In particular, stochastic inflation [8–11] (or, in general, for a semiclassical treatment for the inflaton field during inflation) has been subject of great interest in the last years. However, one of the problems with this approach is that one must to make a perturbative expansion of the scalar field potential in terms of the quantum fluctuations of the inflaton field to finally give a solution for a first order expansion in the equation of motion for these fluctuations [12]. This is a good approximation because such that fluctuations are small during inflation on cosmological scales. However, the predictions of the inflationary theory

*E-mail address: mbellini@mdp.edu.ar

could be significantly improved by using a nonperturbative calculation for the inflaton field φ . Of course, it is impossible to make from a 4D quantum field formalism, but could be developed from a scalar quantum (inflaton) field in a 5D vacuum state with a purely kinetic density Lagrangian. The aim of this work consists to develop a nonperturbative scalar quantum field theory from a 5D apparent vacuum state defined as a purely kinetic 5D density Lagrangian of a scalar field minimally coupled to gravity in a 5D Ricci-flat canonical metric [13]. To make it, we consider the 5D canonical metric [14]

$$dS^2 = \psi^2 dN^2 - \psi^2 e^{2N} dr^2 - d\psi^2, \quad (1)$$

where $dr^2 = dx^2 + dy^2 + dz^2$. Here, the coordinates $(N, \vec{r})$ are dimensionless and the fifth coordinate ψ has spatial unities. We shall assume in what follows that the extra dimension is spacelike and that the universe is 3D spatially flat, isotropic and homogeneous. The metric (1) describes a flat 5D manifold in apparent vacuum ($G_{AB} = 0$). We consider a diagonal metric because we are dealing only with gravitational effects, which are the important ones during inflation. To describe neutral matter in a 5D geometrical vacuum (1) we can consider the Lagrangian

$${}^{(5)}L(\varphi, \varphi_{,A}) = -\sqrt{\left|\frac{{}^{(5)}g}{{}^{(5)}g_0}\right|} {}^{(5)}\mathcal{L}(\varphi, \varphi_{,A}), \quad (2)$$

where $|{}^{(5)}g| = \psi^8 e^{6N}$ is the absolute value of the determinant for the 5D metric tensor with components g_{AB} (A, B take the values 0, 1, 2, 3, 4) and $|{}^{(5)}g_0| = \psi_0^8 e^{6N_0}$ is a constant of dimensionalization determined by $|{}^{(5)}g|$ evaluated at $\psi = \psi_0$ and $N = N_0$. In this work we shall consider $N_0 = 0$, so that ${}^{(5)}g_0 = \psi_0^8$. Here, the index “0” denotes the values at the end of inflation. Furthermore, we shall consider an action

$$I = -\int d^4x d\psi \sqrt{\left|\frac{{}^{(5)}g}{{}^{(5)}g_0}\right|} \left[\frac{{}^{(5)}R}{16\pi G} + \mathcal{L}(\varphi, \varphi_{,A}) \right],$$

where φ is a scalar field minimally coupled to gravity and G is the gravitational constant. Furthermore, ${}^{(5)}R$ is the 5D Ricci scalar, which of course, is zero for the 5D flat metric (1).

Since the 5D metric (1) describes a manifold in apparent vacuum, the density Lagrangian $\mathcal{L}$ in (2) must to be

$${}^{(5)}\mathcal{L}(\varphi, \varphi_{,A}) = \frac{1}{2} g^{AB} \varphi_{,A} \varphi_{,B}, \quad (3)$$

which represents a free scalar field. In other words, we define the vacuum as a purely kinetic 5D-lagrangian on a globally 5D-flat metric [in our case, the metric (1)]. To describe the metric in physical coordinates we can make the following transformations:

$$t = \psi_0 N, \quad R = \psi_0 r, \quad \psi = \psi, \quad (4)$$

such that we obtain the 5D metric

$$dS^2 = \left(\frac{\psi}{\psi_0}\right)^2 \left[dt^2 - e^{2t/\psi_0} dR^2 \right] - d\psi^2, \quad (5)$$

where t is the cosmic time and $R^2 = X^2 + Y^2 + Z^2$. This metric is the Ponce de Leon one [15], and describes a 3D spatially flat, isotropic and homogeneous extended (to 5D) FRW metric in a de Sitter expansion [4].

II. QUANTUM FIELD THEORY IN A 5D APPARENT VACUUM

Taking into account the metric (1) and the Lagrangian (2), we obtain the equation of motion for φ

$$\left(2\psi \frac{\partial \psi}{\partial N} + 3\psi^2\right) \frac{\partial \varphi}{\partial N} + \psi^2 \frac{\partial^2 \varphi}{\partial N^2} - \psi^2 e^{-2N} \nabla_r^2 \varphi - 4\psi^3 \frac{\partial \varphi}{\partial \psi} - 3\psi^4 \frac{\partial N}{\partial \psi} \frac{\partial \varphi}{\partial \psi} - \psi^4 \frac{\partial^2 \varphi}{\partial \psi^2} = 0, \quad (6)$$

where $\frac{\partial N}{\partial \psi}$ is zero because the coordinates $(N, \vec{r}, \psi)$ are independents. Hence, we obtain

$$\overset{\star\star}{\varphi} + 3 \overset{\star}{\varphi} - e^{-2N} \nabla_r^2 \varphi - \left[4\psi \frac{\partial \varphi}{\partial \psi} + \psi^2 \frac{\partial^2 \varphi}{\partial \psi^2}\right] = 0, \quad (7)$$

where the overstar denotes the derivative with respect to N and $\varphi \equiv \varphi(N, \vec{R}, \psi)$. To simplify its structure we can make the transformation $\varphi = \chi e^{-3N/2} \left(\frac{\psi_0}{\psi}\right)^2$, so that we obtain the 5D generalized Klein-Gordon like equation for the redefined field $\chi(N, \vec{r}, \psi)$:

$$\overset{\star\star}{\chi} - \left[e^{-2N} \nabla_r^2 + \left(\psi^2 \frac{\partial^2}{\partial \psi^2} + \frac{1}{4}\right)\right] \chi = 0. \quad (8)$$

The field χ can be written in terms of a Fourier expansion

$$\chi(N, \vec{r}, \vec{\psi}) = \frac{1}{(2\pi)^{3/2}} \int d^3 k_r \int d k_\psi \left[a_{k_r k_\psi} e^{i(\vec{k}_r \cdot \vec{r} + \vec{k}_\psi \cdot \vec{\psi})} \xi_{k_r k_\psi}(N, \psi) + a_{k_r k_\psi}^\dagger e^{-i(\vec{k}_r \cdot \vec{r} + \vec{k}_\psi \cdot \vec{\psi})} \xi_{k_r k_\psi}^*(N, \psi) \right], \quad (9)$$

where the asterisk denotes the complex conjugate and $(a_{k_r k_\psi}^\dagger, a_{k_r k_\psi})$ are the creation and annihilation operators such that

$$\left[a_{k_r k_\psi}, a_{k'_r k'_\psi}^\dagger \right] = \delta^{(3)}(\vec{k}_r - \vec{k}'_r) \delta(\vec{k}_\psi - \vec{k}'_\psi). \quad (10)$$

$$\left[a_{k_r k_\psi}^\dagger, a_{k'_r k'_\psi}^\dagger \right] = \left[a_{k_r k_\psi}, a_{k'_r k'_\psi} \right] = 0. \quad (11)$$

Furthermore, the commutation relation between χ and $\overset{\star}{\chi}$ is

$$\left[\chi(N, \vec{r}, \psi), \overset{\star}{\chi}(N, \vec{r}', \psi') \right] = i \delta^{(3)}(\vec{r} - \vec{r}') \delta(\vec{\psi} - \vec{\psi}'). \quad (12)$$

In order to the commutation equation (12) holds the following renormalization condition must to be fulfilled:

$$\xi_{k_r k_\psi} \left(\overset{\star}{\xi}_{k_r k_\psi} \right)^* - \left(\xi_{k_r k_\psi} \right)^* \overset{\star}{\xi}_{k_r k_\psi} = i. \quad (13)$$

Hence, the equation for the modes $\xi_{k_r k_\psi}(N, \psi)$ that complies with the condition (13) in a 4D de Sitter expansion will be

$$\overset{\star\star}{\xi}_{k_r k_\psi} + \left[e^{-2N} k_r^2 - \left(\frac{1}{4} - \psi^2 k_\psi^2 \right) \right] \xi_{k_r k_\psi} = 0. \quad (14)$$

The general solution for this equation is

$$\xi_{k_r k_\psi}(N, \psi) = G_1(\psi) \mathcal{H}_\nu^{(1)}[k_r e^{-N}] + G_2(\psi) \mathcal{H}_\nu^{(2)}[k_r e^{-N}], \quad (15)$$

where $\mathcal{H}_\nu^{(1,2)}[x] = \mathcal{J}_\nu[x] \pm i\mathcal{Y}_\nu[x]$ are the Hankel functions, $\mathcal{J}_\nu[x]$ and $\mathcal{Y}_\nu[x]$ are the first and second kind Bessel functions, and $\nu = \frac{\sqrt{1-4k_\psi^2\psi^2}}{2}$. Furthermore, the functions $G_1(\psi)$ and $G_2(\psi)$ are arbitrary functions constrained by the renormalization condition (13)

$$[G_1(\psi) - G_2(\psi)][G_1(\psi) + G_2(\psi)] = \frac{\pi}{4}. \quad (16)$$

In this paper we shall choose the generalized Bunch-Davis vacuum: $G_1(\psi) = 0$ and $G_2(\psi) = i\frac{\sqrt{\pi}}{2}$. The squared φ fluctuations are given by

$$\langle \varphi^2 \rangle = \frac{1}{(2\pi)^3} \int d^3 k_r \int dk_\psi \xi_{k_r k_\psi} \xi_{k_r k_\psi}^*. \quad (17)$$

A k_r -scale invariant power spectrum results from $\nu = 3/2$, for which

$$ik_\psi = -\frac{\sqrt{2}}{\psi}. \quad (18)$$

III. 4D DE SITTER EXPANSION

We can take a foliation $\psi = \psi_0$ in the metric (5), such that the effective 4D metric results

$$dS^2 \rightarrow ds^2 = dt^2 - e^{2t/\psi_0} dR^2, \quad (19)$$

which describes 4D globally isotropic and homogeneous expansion of a 3D spatially flat, isotropic and homogeneous universe that expands with a Hubble parameter $H = 1/\psi_0$ (in our case a constant) and a 4D scalar curvature ${}^{(4)}\mathcal{R} = 6(\dot{H} + 2H^2)$. Note that in this particular case the Hubble parameter is constant so that $\dot{H} = 0$.

The 4D energy density ρ and the pressure p are [14]

$$8\pi G \langle \rho \rangle = 3H^2, \quad (20)$$

$$8\pi G \langle p \rangle = -3H^2, \quad (21)$$

where $G = M_p^{-2}$ is the gravitational constant and $M_p = 1.2 \cdot 10^{19} \text{ GeV}$ is the Planckian mass. Furthermore, the universe describes a vacuum equation of state: $p = -\rho$, such that

$$\langle \rho \rangle = \left\langle \frac{\dot{\varphi}^2}{2} + \frac{a_0^2}{2a^2} (\vec{\nabla} \varphi)^2 + V(\varphi) \right\rangle, \quad (22)$$

where the brackets denote the 4D expectation vacuum and the cosmological constant Λ gives the vacuum energy density $\langle \rho \rangle = \frac{\Lambda}{8\pi G}$. Thus, Λ is related with the fifth coordinate by means of $\Lambda = 3/\psi_0^2$ [15]. Furthermore, the 4D Lagrangian is given by

$${}^{(4)}\mathcal{L}(\varphi, \varphi_{,\mu}) = -\sqrt{\left|\frac{{}^{(4)}g}{{}^{(4)}g_0}\right|} \left[\frac{1}{2} g^{\mu\nu} \varphi_{,\mu} \varphi_{,\nu} + V(\varphi) \right], \quad (23)$$

where the effective potential for the 4D FRW metric [13], is

$$V(\varphi) = -\frac{1}{2} g^{\psi\psi} \varphi_{,\psi} \varphi_{,\psi} \Big|_{\psi=\psi_0} = \frac{1}{2} \left(\frac{\partial \varphi}{\partial \psi} \right)^2 \Big|_{\psi=\psi_0} \quad (24)$$

In our case this potential takes the form

$$V(\varphi) = \left(\frac{2}{\psi_0^2} - \frac{k_{\psi_0}^2}{2} - \frac{2ik_{\psi_0}}{\psi_0} \right) \varphi^2(t, \vec{R}, \psi_0), \quad (25)$$

where k_{ψ_0} is the wavenumber for $\psi = \psi_0$. Furthermore the effective 4D motion equation for φ is

$$\ddot{\varphi} + \frac{3}{\psi_0} \dot{\varphi} - e^{-2t/\psi_0} \nabla_R^2 \varphi - \left[4 \frac{\psi}{\psi_0^2} \frac{\partial \varphi}{\partial \psi} + \frac{\psi^2}{\psi_0^2} \frac{\partial^2 \varphi}{\partial \psi^2} \right] \Big|_{\psi=\psi_0} = 0, \quad (26)$$

which means that the effective derivative (with respect to φ) for the potential, is

$$V'(\varphi)|_{\psi=\psi_0} = \alpha \varphi(\vec{R}, t, \psi_0), \quad (27)$$

with

$$\alpha = \frac{2}{\psi_0^2} + k_{\psi_0}^2. \quad (28)$$

Now we can make the following transformation:

$$\varphi(\vec{R}, t) = e^{-\frac{3t}{2\psi_0}} \chi(\vec{R}, t). \quad (29)$$

Note that now $\varphi \equiv \varphi(\vec{R} = \psi_0 r, t = \psi_0 N, \psi = \psi_0) = e^{-3t/(2\psi_0)} \chi(\vec{R}, t)$, where [see eq. (9)] $\chi(\vec{R}, t) = \chi(\vec{R} = \psi_0 r, t = \psi_0 N, \psi = \psi_0)$:

$$\chi(\vec{R}, t) = \frac{1}{(2\pi)^{3/2}} \int d^3 k_R \int dk_\psi \left[a_{k_R k_\psi} e^{i(\vec{k}_R \cdot \vec{R} + k_\psi \cdot \vec{\psi})} \xi_{k_R k_\psi}(t, \psi) + c.c. \right] \delta(k_\psi - k_{\psi_0}). \quad (30)$$

Hence, we obtain the following 4D Klein-Gordon equation for χ

$$\ddot{\chi} - \left[e^{-\frac{2t}{\psi_0}} \nabla_R^2 + \frac{9}{4\psi_0^2} - \alpha \right] \chi = 0. \quad (31)$$

The equation of motion for the time dependent modes $\xi_{k_R k_{\psi_0}}(t)$ will be

$$\ddot{\xi}_{k_R k_{\psi_0}} + \left[k_R^2 e^{-\frac{2t}{\psi_0}} - \left(\frac{9}{4\psi_0^2} - \alpha \right) \right] \xi_{k_R k_{\psi_0}} = 0, \quad (32)$$

where the effective squared mass μ^2 of the modes $\xi_{k_R k_{\psi_0}}$ is given by

$$\mu^2 = \frac{9}{4\psi_0^2} - \alpha. \quad (33)$$

Here, α describes the self-interaction squared mass of the redefined inflaton field χ due to the expansion of the universe, and the term $\frac{9}{4\psi_0^2}$ represents its bare squared mass due to the expansion. Note that both terms in (33) has a geometrical origin because they are induced by the fifth coordinate.

The general solution of eq. (32) is

$$\xi_{k_R k_{\psi_0}} = G_1(\psi_0) \mathcal{H}_{\frac{3-n_s}{2}}^{(1)} [k_R \psi_0 e^{-t/\psi_0}] + G_2(\psi_0) \mathcal{H}_{\frac{3-n_s}{2}}^{(2)} [k_R \psi_0 e^{-t/\psi_0}]. \quad (34)$$

Here, $\frac{(3-n_s)}{2} = \sqrt{\frac{9}{4} - (2 + k_{\psi_0}^2 \psi_0^2)}$ and n_s is the spectral index of the $\langle \varphi^2 \rangle$ -spectrum on super Hubble (SH) scales when it is considered the Bunch-Davis vacuum: $G_1(\psi_0) = 0$, $G_2(\psi_0) = i\sqrt{\pi}/2$

$$\langle \varphi^2 \rangle|^{SH} = \int_{k_*}^{\epsilon k_0} \frac{dk}{k} \mathcal{P}(t, k) \simeq A(t) \int_{k_*}^{\epsilon k_0} \frac{dk}{k} k^{n_s}, \quad (35)$$

where $A(t)$ is a time dependent function, $k_0(t) = \frac{e^{t/\psi_0}}{\psi_0} \sqrt{\frac{9}{4} - \alpha \psi_0^2}$, $\epsilon \ll 1$, $\mathcal{P}(t, k)$ is the power of the spectrum and k_* is the absolute value for the wave vector related to the physical wavelength at the moment the horizon entry. This maximal physical scale is supported by causal arguments. Note that this result corresponds to an effective self-interaction squared mass α for the inflaton field

$$\alpha = 2H^2 + k_{\psi_0}^2; \quad \text{for } \psi_0 = 1/H, \quad (36)$$

when the standard semiclassical approach in a de Sitter expansion is considered [12]. For $n_s \ll 1$, it can be approximated to $n_s \simeq \frac{2(2+k_{\psi_0}^2 \psi_0^2)}{3}$, where

$$ik_{\psi_0} \simeq -\frac{\sqrt{2-3n_s/2}}{\psi_0} = -\sqrt{2-3n_s/2} H. \quad (37)$$

It is well known that the universe has a power spectrum which is very close to a scale invariant one on cosmological scales [16], so that the equation (37) is a good approximation. In such that case we obtain the following valued expressions for $V(\varphi)$ and $V'(\varphi)$:

$$V(\varphi)|_{(\psi=\psi_0, n_s=0)} = \frac{(4+2\sqrt{2})}{\psi_0^2} \varphi^2(\vec{R}, t, \psi_0), \quad (38)$$

$$V'(\varphi)|_{(\psi=\psi_0, n_s=0)} = 0. \quad (39)$$

where the equation (39) corresponds to $\alpha = 0$ in (31). Therefore, α becomes zero for a super Hubble scale invariant power spectrum of $\langle \varphi^2 \rangle$. Note that this result disagrees with the result obtained using a semiclassical 4D treatment for the inflaton field in a de Sitter expansion [12]. Furthermore, we can see from eq. (39) that the effective 4D parameter of mass for φ

$$m_{eff}^2 = \frac{2(4+2\sqrt{2})}{\psi_0^2}, \quad (40)$$

is nonzero. It describes the inflaton φ -squared mass geometrically induced by the fifth coordinate ψ on the hypersurface $\psi = \psi_0$, when the self-interaction is absent in a de Sitter expansion: $\alpha = 0$. It is easy to see that $m_{eff}^2 > \mu^2|_{\alpha=0}$. However, both masses have a different origin, because μ^2 is the squared mass related to each $\xi_{k_R k_{\psi_0}}$ mode of χ and m_{eff}^2 is the effective squared mass of the nonperturbative field $\varphi(\vec{R}, t, \psi_0)$, with back-reaction effects included.

A. Energy density fluctuations

Once the time dependent modes $\xi_{k_R k_{\psi_0}}$ are known, we can obtain the effective 4D expectation value for the energy density

$$\begin{aligned} \langle \rho \rangle_{\psi=\psi_0} &= \frac{e^{-3t/\psi_0}}{(2\pi)^3} \int_{k_*}^{k_p} d^3 k_R \left\{ \left[\frac{(26 + (3 - n_s)^2)}{8\psi_0^2} - \frac{2ik_{\psi_0}}{\psi_0} + \frac{e^{-2t/\psi_0} k_R^2}{2} \right] \xi_{k_R k_{\psi_0}} \xi_{k_R k_{\psi_0}}^* \right. \\ &\quad \left. + \frac{1}{2} \dot{\xi}_{k_R k_{\psi_0}} \dot{\xi}_{k_R k_{\psi_0}}^* - \frac{3}{4\psi_0} (\xi_{k_R k_{\psi_0}} \dot{\xi}_{k_R k_{\psi_0}}^* + \dot{\xi}_{k_R k_{\psi_0}} \xi_{k_R k_{\psi_0}}^*) \right\}, \end{aligned} \quad (41)$$

where $k_p = G^{-1/2}$ is the absolute value of the Planckian wave vector and k_* is the inverse of the Hubble's radius at the moment ($t = t_*$) the horizon entry.

On the other hand, if we define $\langle \rho \rangle_{\psi=\psi_0}^{(0)}$ the k_R -zero mode 4D expectation value for the energy density

$$\begin{aligned} \langle \rho \rangle_{\psi=\psi_0}^{(0)} &= \frac{e^{-3t/\psi_0}}{(2\pi)^3} \int_{k_*}^{k_p} d^3 k_R \left\{ \left[\frac{(26 + (3 - n_s)^2)}{8\psi_0^2} - \frac{2ik_{\psi_0}}{\psi_0} \right] \xi_{0k_{\psi_0}} \xi_{0k_{\psi_0}}^* \right. \\ &\quad \left. + \frac{1}{2} \dot{\xi}_{0k_{\psi_0}} \dot{\xi}_{0k_{\psi_0}}^* - \frac{3}{4\psi_0} (\xi_{0k_{\psi_0}} \dot{\xi}_{0k_{\psi_0}}^* + \dot{\xi}_{0k_{\psi_0}} \xi_{0k_{\psi_0}}^*) \right\}, \end{aligned} \quad (42)$$

on the effective 4D spatially isotropic and homogeneous (de Sitter) FRW metric (19), hence the energy density fluctuations (on arbitrary scales) will be

$$\delta\rho/\rho = \frac{\langle \rho \rangle_{\psi=\psi_0} - \langle \rho \rangle_{\psi=\psi_0}^{(0)}}{\langle \rho \rangle_{\psi=\psi_0}}. \quad (43)$$

B. Super Hubble energy density fluctuations

It is well known that the universe is nearly scale invariant on cosmological scales [16]: $|n_s| \ll 1$. We shall consider this range for the spectrum on the range $k_* < k_R < \epsilon k_0(t)$. In order to compute the SH density energy fluctuations we can remember the small-argument limit for the second kind Hankel function: $\mathcal{H}_\nu^{(2)}[x] \simeq -\frac{i}{\pi} \Gamma(\nu) (x/2)^{-\nu}$, for $\nu > 0$, which in our case is valid to describe the super Hubble (cosmological) φ -field time dependent modes $\xi_{k_R k_{\psi_0}}$ in eq. (34). With this approach, we obtain

$$\xi_{k_R k_{\psi_0}} \xi_{k_R k_{\psi_0}}^* \simeq \frac{2^{1-n_s}}{\pi} \Gamma^2 \left(\frac{3-n_s}{2} \right) [k_R \psi_0 e^{-t/\psi_0}]^{-(3-n_s)}, \quad (44)$$

$$\dot{\xi}_{k_R k_{\psi_0}} \dot{\xi}_{k_R k_{\psi_0}}^* \simeq \left(\frac{3-n_s}{2} \right)^2 \frac{\Gamma^2 \left(\frac{3-n_s}{2} \right)}{2^{1+n_s} \pi \psi_0^2} [k_R \psi_0 e^{-t/\psi_0}]^{-(1-n_s)}, \quad (45)$$

$$\frac{\xi_{k_R k_{\psi_0}} \dot{\xi}_{k_R k_{\psi_0}}^* + \dot{\xi}_{k_R k_{\psi_0}} \xi_{k_R k_{\psi_0}}^*}{2} \simeq \left(\frac{3-n_s}{2} \right) \frac{\Gamma^2 \left(\frac{3-n_s}{2} \right)}{2^{n_s} \pi \psi_0} [k_R \psi_0 e^{-t/\psi_0}]^{-(2-n_s)}, \quad (46)$$

and for the zero-modes, we obtain

$$\xi_{0k_{\psi_0}} \xi_{0k_{\psi_0}}^* \simeq \frac{2}{\pi} \Gamma^2 \left(\frac{3}{2} \right) [k_R \psi_0 e^{-t/\psi_0}]^{-3}, \quad (47)$$

$$\dot{\xi}_{0k_{\psi_0}} \dot{\xi}_{0k_{\psi_0}}^* \simeq \frac{9}{8\pi} \frac{\Gamma^2 \left(\frac{3}{2} \right)}{\psi_0^2} [k_R \psi_0 e^{-t/\psi_0}]^{-1}, \quad (48)$$

$$\frac{\xi_{0k_{\psi_0}} \dot{\xi}_{0k_{\psi_0}}^* + \dot{\xi}_{0k_{\psi_0}} \xi_{0k_{\psi_0}}^*}{2} \simeq \frac{3}{2\pi} \frac{\Gamma^2 \left(\frac{3}{2} \right)}{\psi_0} [k_R \psi_0 e^{-t/\psi_0}]^{-2}. \quad (49)$$

Hence, on cosmological scales (i.e., for $k^2 \ll k_0^2$), the late times energy density fluctuations renormalized at $n_s = 0$ will be approximately (i.e., at first order in n_s)

$$\left. \frac{\delta \rho}{\rho} \right|_{t=t_*}^{SH} \simeq n_s \ln [k_R]_{k_*}^{\epsilon_{k_0}(t_*)}, \quad (50)$$

where t_* is the time when the horizon entry and we have made use of the expression (37). Note that eq. (50) describes the departure of $\delta \rho / \rho$ on super Hubble scales with respect to scale invariant density fluctuations ($n_s = 0$). Furthermore, the expression (50) is consistent for

$$n_s \ln [k_R]_{k_*}^{\epsilon_{k_0}(t_*)} < 1. \quad (51)$$

Hence, if $k_* e^{-H_* t_*} = H_*$ is the Hubble parameter when the horizon entry and we consider $\frac{\epsilon \sqrt{\frac{9}{4} - \alpha \psi_0^2}}{H_* \psi_0} e^{(\psi_0^{-1} - H_*) t_*} > e^{60}$, one obtains the following inequality¹:

$$60 < \ln [k_R]_{k_*}^{\epsilon_{k_0}(t_*)} < \frac{1}{n_s}, \quad (52)$$

which is fulfilled for

$$n_s < 1/60. \quad (53)$$

This is an important result which is in good agreement with observation [17].

¹In the calculations we are considering $\psi_0^{-1} = H$, in consistence with the 4D FRW metric (19)

IV. FINAL COMMENTS

We have developed a 5D nonperturbative quantum scalar field formalism and studied the particular case of a 4D de Sitter inflationary expansion in the framework of the induced-matter theory developed by Wesson and co-workers. The results are very interesting, because we note some differences with respect to those obtained by means of the semiclassical expansion in a de Sitter expansion for the universe. 1) First all, the effective scalar 4D potential appears to be quadratic in φ (see eq. (25), meanwhile in the standard semiclassical treatment is a constant V_0 (see for example [12]). 2) The spectral index $n_s < 1/60$ depends on the self-interaction squared mass α (which should be very small if the slow-roll conditions hold), but not on the mass of the inflaton field m_{eff} (as in the standard semiclassical approach to inflation). Notice this result suggests that $k_{\psi_0} \simeq i\sqrt{2}/\psi_0 = i\sqrt{2}H$. 3) Furthermore, the Hubble parameter in a de Sitter expansion (as in the semiclassical approach to inflation) is a constant, but in our case, its value being given by the inverse of the fifth coordinate $H = 1/\psi_0$ for a foliation $\psi = \psi_0$ of the 5D Ricci-flat metric (5).

Finally, the nonperturbative treatment for φ here developed can be extended to other (more realistic) inflationary models with time dependent Hubble parameters and, more generally, to cosmological models dynamically governed by scalar fields.

M.B. Acknowledges CONICET, AGENCIA and UNMdP (Argentina) for financial support.

REFERENCES

- [1] N. Akdani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429**, 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436**, 257 (1998).
- [2] D. Youm, Phys. Rev. **D62**, 084002 (2000); R. Maartens, Phys. Rev. **D62**, 084023 (2000); A. Chamblin, Class. Quantum Grav. **18**, L-17 (2001); P. S. Wesson, B. Mashhoon, H. Liu, W. N. Sajko, Phys. Lett. **B456**, 34 (2001); J. Ponce de Leon, Phys. Lett. **B523**, 311 (2001).
- [3] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
- [4] For a review of STM theory the reader can see J. M. Overduin and P. S. Wesson, Phys. Rep. **283**, 303 (1997).
- [5] B. Mashhoon, H. Liu and P. S. Wesson, Phys. Lett. **B331**, 305 (1994).
- [6] B. Mashhoon and P. Wesson, Class. Quant. Grav. **21**, 3611 (2004).
- [7] D. H. Lyth and A. Riotto, Phys. Rept. **314**, 1 (1999).
- [8] S. Habib, Phys. Rev. **D46**, 2408 (1992).
- [9] M. Mijic, Phys. Rev. **D49**, 6434 (1994).
- [10] E. J. Copeland, E. W. Colb, A. R. Liddle and J. E. Lidsay, Phys. Rev. **D48**, 2529 (1993).
- [11] M. Liguori, S. Matarrese, M. Musso and A. Riotto, JCAP 0408: 011 (2004).
- [12] M. Bellini, H. Casini, R. Montemayor and P. Sisterna, Phys. Rev. **D54**, 7172 (1996).
- [13] J. E. Madriz Aguilar and M. Bellini, Phys. Lett. **B596**, 116 (2004); J. E. Madriz Aguilar and M. Bellini, Eur. Phys. J. **C38**, 367 (2004).
- [14] D. S. Ledesma and M. Bellini, Phys. Lett. **B581**, 1 (2004).
- [15] J. Ponce de Leon, Gen. Rel. Grav. **20**, 539 (1988).
- [16] A. H. Jaffe et. al, Phys. Rev. Lett. **86**, 3475 (2001).
- [17] Review of Particle Physics: Phys. Lett. **B592**, 207 (2004).